

Bibliography

- Abdel-adar, K., Myaskovsky, L., Kapro, I., Shah, J., Hess, R., Dew, M.A., & Unruh, M. (2009). Individual quality of life in chronic kidney disease : Influence of age and dialysis modality. *Clinical Journal of American Society Nephrology*;4(4):711-18 Epub 2009,April2009.
- Abdel-Khader, K., Unruh, M.L., & Weishbord, S.D. (2009). Symptom burden, depression and quality of life in chronic and end stage kidney disease. *Cl. J. Am. Soc. Nephrol.*, 4(6), 1057 to 1064.
- American Nurses Association (1995). Nursing a social policy statement. Washington D C
- American Psychiatric Association (1994). – *Diagnostic and statistical manual of mental disorder* (4th ed.). Washington D C.
- American Psychological Association Publication (2002). *Manual of the American Psychological Association* (5th ed.). Washington D C
- ANNA position statement: Chronic kidney disease (2007) American Nephrology Nurses Association. Available from <http://www.annanurse.org>
- Agarwal, R., Alborzi, P., Satyan, S., & Light, R. P. (2009). Dry weight reduction in hypertensive - hemodialysis patients (DRIP): a randomized, controlled trial. *Hypertension*, 53(3), 500-7. Epub 2009 Jan 19.
- Agarwal R. (2002) Assessment of blood pressure in hemodialysis patients - *semin. Dial.* 15(5), 299-304.
- Agarwal, S. K., Dash, S. C., & Irshad, M., (2005). Prevalance of chronic failure in adults in Delhi, India, *Nephro Dial Transplant*,20, 1638-1642.
- Agarwal, S K., & Srivastava, R. K. (2009). Chronic kidney disease in India: Challenges and solutions. *Nephron Clinical Practice*;111(3):197-200.
- Ageborg, M., Allenius, B.L., & Oderijaii, O. (2005). Quality of life, self-care ability and sense of coherence in hemodialysis patients: A comparative study. *Hemodial. Int.*, 1, 58-14.
- Ahari, S.S., Bashardoost, B., Amani, F., Mohammadi, M.A., Siahpoosh, H., & Arab, R. (2007). Quality of life in hemodialysis patients at Ardebil University of Medical Sciences (Arums) and factors affecting it. *Journal of Biological Sciences*, 2(5), 529-533.
- Alligood, M.R., & Tomey A.M. (2006). *Nursing Theory Utilization and Application*, (3rd ed.). USA., Mosby pp255-276.

American Kidney Fund – www.akfinc.org.

- Anderson, R.M., & Funnell, M.M. (2005). Patient empowerment: Reflections on the challenge of fostering the adoption of a new paradigm. *Patient Education and Counseling*, 57, 153-157.
- Arora, P., & Verrelli, M., (2009). - *Chronic Renal Failure: e-Medicine Nephrology*. <http://emedicine.medscape.com>. Retrieved on 08.01.09.
- Bolander,V.B (1994) *Sorenson and Luckmann's Basic Nursing: A psycho physiologic approach* (3rded.). Philadelphia, W. B. Saunders.
- Brachy, L. (2009). Comprehensive dialysis study provides data on longevity health status in ESRD. Medscape Medical News. *Cl. J. Ans. Soc. Nephrology*, 4, 645-650. Available <http://www.medscape.com/> resume/end stage - renal disease.
- Bots, C.P., Brad, H.S., Veerman, E.C., & I., Korevaan, J.C. (2005). - Chewing gum and a saliva substitute alleviate thirst and xerostmia in patients on hemodialysis. *Nephrology Dialysis Transplantation*, 20, 578-584.
- Barebdse, S.M. (2005). The Renal Treatment Satisfaction Questionnaire; A measure of satisfaction with treatment for chronic kidney failure. *American Journal of Kidney Diseases*, 45(3), 572-579.
- Buysse Reybolds, & Monk Burman Kuph. (1989). The Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. *Psychiatry Research*, 28, 193-213.
- Burrows, H. (2005). Chronic kidney disease: An overview. *American Journal of Nursing*, 2, 40-50
- Barclay, L. (2009). Patients with less dialysis knowledge may be less likely to use permanent arteriovenous access. *Clinical Journal of the American Society of Nephrology*, 4.
- Burrowes, J. D., & Ramer, N.J. (2006) Removal of potassium from tuberous root vegetables by teaching. *Journal of Renal Nutrition: the official journal of the Council of Renal Nutrition of the National Kidney Foundation*, 16(4), 304-311.
- Barnet, T., Li Yoong, T., Pinikahana, J., & Si-Yen, T., (2008). Fluid compliance among patients having hemodialysis can an educational program made a difference? *J. Adv. Nurs.* 61(3), 300-6.
- Barakzoy, A.S., & Moss, A.H. (2009). Efficacy of the World Health Organization analgesic ladder to treat pain in end-stage renal disease. *Clin. J. Am. Soc. Nephrol.*, 4(3), 517.
- Bakris, G.L., & Ritz, E. (2009). World Kidney Day Steering Committee, The message for World Kidney Day 2009: Hypertension and kidney disease, A marriage that should be prevented, *Kidney International* 75, 449-452.

- Brown, L., Ruth, L. AIDS Quest: The HIV/AIDS survey library excerpted from "A review of approaches and instruments for assessing health-related quality of life". *Tulane University / Horizons*, www.popcouncil.org
- Batalden, P., Ogrinc, G., & Batalden, M. (2006). From one to many. *Journal of Interprofessional Care*, 20, 549-551.
- Brunelli, S.M., & Berns, J.S. (2009). *Anemia in chronic kidney disease and end-stage renal disease Nephrology Rounds*, 7(8), www.nephrologyrounds.org
- Brenner BM. (1986) *The kidney* (3rd ed.). Philadelphia W B Saunders co.
- Black, M.J., & Hawks, H.J. (2007). Medical surgical nursing Vol.1 7th ed. New Delhi Elsevier.
- Carpentor, L.J. (1993) – *Nursing Diagnosis Application in Clinical Practice*, Philadelphia.
- Cavanaugh, K.L., Wingard, R.L., Hakim, R.H., Elasy, T.A., Ikizler, T.A. (2009). Patient Dialysis Knowledge Is Associated with Permanent Arteriovenous Access Use in Chronic Hemodialysis, *Clin J Am Soc Nephrol* 4: 950-956
- Cole, C., & Kathy. (2007). Sleep Disruption in older Adults: Harmful and by no means inevitable, it should be assessed for and treated, *American Journal of Nursing*, 107(5), 40-49.
- Conn VS, Halfdahal AR, Brown SA, Brown LM. (2008). Meta-Analysis of patient education intervention to increase physical activity among chronically adults *Patient Edu Couns*, 70(2),157-172online 2007 Nov 26 doi:10.1016/J.PEC2007.10.004
- Carreon, M., Fried, F.L., Palevsky, P.M., & Kimmel, P.L. (2008). Clinical correlates treatment of bone/joint pain and difficulty with sexual arousal in patients on maintenance hemodialysis. *Hemodialysis International*, 12(2), 268-274.
- Close, A. (1988). Patient Education: a literature review. *Journal of advanced Nursing*, 13(2), 2003-13.
- Curtin, R.B., Johnson. H.K. & Schatell, D. (2004). The Peritoneal Dialysis Experience: Insight from Long Term Patients. *Nephrology Nursing Journal*, 31(6), 615-625.
- Cummings, G.G. (2006). Development and validation of a desired measure of research utilization by nurses. *Nursing Research*, 55(3), 149-160.
- Ching, L.Y., Luan, T.S., & Jyy, W.T. (2007). Intervention of quality of life group in hemodialysis patients. *Journal of Clinical Nursing*, 16(Supplement C), 268-275.
- Castner, D., & Douglas, C. (2005). *Chronic Kidney Disease Nurisng*, 35(12), 58-64.

- Campoy, S. (2005). Pharmacology and chronic kidney disease, *American Journal of Nursing*, 105(9), 60-72.
- Cohen, S.D., Patel, S.S., Khetpal, P., Peterson, R.A., & Kimmel, P.L. (2007). Pain, sleep disturbance and quality of life in patients with chronic kidney disease. *Clin J. Am. Soc. Nephrol*, 2, 919-925.
- Chug, K.S. (1994). Three decades of nephrology, *J. Postgrad. Med.* (Serial online) (cited 2005, June 6); 40, 103-108 available from [http://www.jgmonline.com/article.asp?assn=0022-3859, 40\(3\), 103, epage=8; aulast=Chug](http://www.jgmonline.com/article.asp?assn=0022-3859, 40(3), 103, epage=8; aulast=Chug).
- Chaturvedi, S., Pant, M., & Yadav, N.G. (2007). Hypertension in Delhi; Prevalence, awareness, treatment and control. *Tropical Doctor*, 37, 142-145 doi:10.1258/004947507781524593.
- Culleton B.F, Walsh M, Klarenbach, S.W., Mortis, G., Scott-Douglas, N., Quinn, R.R., Tonelli, M., Donnelly, S., Friedrich, M.G., Kumar, A., Mahallati, H., Hemmelgarn, B.R., & Manns, B.J. (2007). Systematic review of the effect of nocturnal hemodialysis on blood pressure, left ventricular hypertrophy, anemia, mineral metabolism, and health-related quality of life. *JAMA* 298(11), 1291–1299.
- Cooper, B.A., Branley, P., Bulfone, L., Collins, J.F., Craig, J.C., Dempster, J., Fraenkel, M.B., Harris, A., Harris, D.C., Johnson, D.W., Kesselhut, J., Luxton, G., Pilmore, A., Pollock, C.A., & Tiller, D.J. (2004). The Initiating Dialysis Early and Late (IDEAL) study: study rationale and design. *Perit Dial Int.*, 24(2), 176-81.
- Carrero, J.J., Qureshi, A.R., Axelsson, J., Avesani, C.M., Suliman, M.E., Kato, S., Bárány, P., Jónsdóttir, S.S., Alvestrand, A., Heimbürger, O., Lindholm, B., & Stenvinkel, P. (2007). Comparison of nutritional and inflammatory markers in dialysis patients with reduced appetite *American Journal of Clinical Nutrition*, 85(3) 695-701.
- Cohen, E. (2005). The cardio-renal anemia syndrome: A triple threat, *Med Medscape Nephrology, Chronic Kidney Disease Expert Column*.
- Cohen, R, Mount, B.M., Strobel, & M.G., Bui, F. (1995). The McGill quality of Life questionnaire: A measure of quality of life appropriate for people with advanced disease. *Pall Med*, 9, 207–19. Retrieved 08.07.2005.
- Chiang, C.K., Peng, Y.S., Chiang, S.S., Yang, C.S., He, Y.H., Hung, K.U., Wu, K.D., Wu, M.S., Cheng, Fang, C., Tsai, T.J., & Chen, W.Y. (2004). Health-related quality of life of hemodialysis patients in Taiwan: A multicenter study. *Blood Purif*, 22, 490–498 doi: 10.1159/000081730. Retrieved on 04.10.05.
- Como, D.N.(Ed.) (1990). Mosby's Medical Nursing and Allied Nursing Dictionary, (3rd ed.). The C. V. Mosby Company, St. Louis.

- Cukor, D., Coplan, J., Brown, C., Friedman, S., Cromwell-Smith, A., Peterson, R.A., & Kimmel, P.L. (2007). Depression and anxiety in urban hemodialysis patients. *Clin J Am Soc Nephrol*, 2, 484-490. doi: 10.2215/CJN.00040107.
- Curtin, R.B., & Thompson, N. (2000). *The Renal Rehabilitation Report Newsletter Editors*, 8(4), www.lifeoptions.org winter.
- Calderon, J., Morales, L., Liu, H., & Hays, R. (2006). Variation in the readability of items within surveys. doi: 10.1177/1062860605283572. *Am J Med Qual*. 21(1), 49-56.
- Curtin, R.B., & Mapes, D.L. (2001). Health care management strategies of long term dialysis survivors. *Nephrol Nurs J*. 28(4), 385-92; discussion 393-4. PMID:12143460
- Carnwell R. (1997). *Evaluative research methodology in nursing and health care* (1st ed.). New York, Churchill Livingston.
- Cleary, J., & Drebbab, J. (2005). Quality of life of patients on hemodialysis for end stage renal disease. *J. Adv. Nsg.*, 5(16), 577-86.
- Clinical practice guidelines for nutrition in chronic renal failure. K/DOQI, National Kidney Foundation. *Am J Kidney Dis*, 2000. 35(6 Suppl 2), S1-140.
- Chronic kidney disease (2006). A guide to select NKF-KDOQI guidelines and recommendation. www.kidney.org/professionals/kls/pdf/Pharmacist_CPG.pdf.
- Chronic kidney disease guideline. <http://www.renal.org/eGFR/eguide.html>.
- Dunning, G. (2005). The choice application and review of topical treatment for skin condition. *Nursing Times*, 101(4), 55.
- Doorenbos, A., Given, B. Given, C. & Verbitsky, N. (2006). Physical functioning effect of behavioral intervention for symptoms among individuals with cancer. *Nursing Research*, 55(3), 161-171.
- Davita Education. (2009). *Diet & Nutition*, <http://www.davita.com/xml/rss.xml>.
- Doak, C., Doak, L., Root, J. (1996). *Teaching Patients with Low Literacy skills* (2nd ed.). Philadelphia, P.A. J.B. Lipincott.
- Doak, C.C., Leonard G. D., & Jane H. R. (1995). *Teaching Patients with Low Literacy Skills*, (2nd ed.). Philadelphia: J.B. Lippincott Company.
- Dember, L.M. (2002). Critical care issues in the patient with chronic renal failure. *Critical Care Clinics*, 4, 421-436.
- Denhareynck, K., Manhaeve, D., Dobbels, F., Garzoni, D., N'olte, C., & Geest, S.D. (2007). Prevalence and consequences of non-adherence of to hemodialysis regimens. *American Journal of Critical Care*, 16, 222-235.

- de Bie, M.K., van Dam, B., Gaasbeek, A., van Buren, M., van Erven, L., Bax, J.J., Schalij, M.J., Rabelink, T.J., & Jukema, J.W. (2009). The current status of interventions aiming at reducing sudden cardiac death in dialysis patients. *Eur Heart J.* 2009 Jul;30(13):1559-64.
- Dagdeviren, A., & Savaser, S. (2003). Education to reduce potassium levels in adolescent, hemodialysis patients, *EDTNA ERCA J.*, 29(4), 195-7.
- Dash, S.C. & Agarwal, S.K. (2006). - Incidence of chronic kidney disease in India – *Nephrol Dial Transplant.*, 21 (1) 232-33.
- DeOreo, P.B, (1997). Hemodialysis patient-assessed functional health status predicts continued survival, hospitalization, and dialysis-attendance compliance. *Am J Kidney Dis.*, 30(2), 04-12.
- Denhaerynck, K. Manhaeve, D., Dobbels, F., Garzoni, D., Nolte, C., & De Geest, S. (2007). Prevalence and consequences of non-adherence to hemodialysis regimens. *Am J Crit Care*, 16(3), 222-35.
- Degoulet, P., Legrain, M., Reach, I., Acme, F., Devries, C., Rojas, P. & Jacobs, C. (1982). Mortality risk factors in patients treated by chronic hemodialysis. *Nephron* 31:103-110 doi: 10.1159/000/82627.
- Eryavuz, N., Yukesel, S., Acartusk, G., Uslam, I., & Nukmber, (2008). Comparison of sleep Quality between hemodialysis and peritoneal dialysis patients. *International Urology and Nephrology*, 40(3), 785 – 791.
- Eghbali, M., Shahqolian, N., Nazari, F., & Babaee, S. (2009). Comparing problems of patients with chronic renal failure undergoing hemodialysis and peritoneal dialysis referring to medical university's hospitals, *IJNMR*, 14(1), 1-5.
- Elderi, S.J., Pisoni, R.L., Akizawa, T., Fissell, R., Andreucci, V.E., Fukuharas, S., Kurokawa, K., Rayner, H.C., Furniss, A.L., Port, F.K. & Saran, R., (2008). Sleep quality predicts quality of life and mortality risk in haemodialysis patients: results from the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Nephrol Dial Transplant*, 23, 998-1004, doi: 10.1093/ndt/gfm630.
- Effendi, K., Sardi, M. A., Jamaludin, R., Abdillah, D.M. & Hernawati (1995). Applications of Dorethea Orem's self-care deficit nursing theory. Available from, *e-learning.unpad kw:*
- Finkelstein, F.O., Wuerth, D., & Finkelstein, S.H. (2009). Health related quality of life and the chronic kidney disease patient: Challenges for the nephrology community. *Kidney International*, 76,946-952 ,doi:10.1038/ki.2009.307.
- Fukuhara, S., Lopes, A. A., Bragg-Grsham, J.L., Kurokawa, K., Mapes, D.L., Akizawa, T., Bommer, J., Canaud, B.J., Port, F. K., & Held, P.J. (2003). Worldwide Dialysis Outcomes and Practice Patterns Study. (2003). Health related quality of life among dialysis patients in three continents: The Dialysis

- Outcomes and Practices Patterns Study. *Kidney International* 2003 Nov; 64 (5): 1903-10. Retrieved 11.09.2006.
- Fallowfield, L.T. (1997). Quality of life data, *Journal of Rehabilitation Outcome Measurement*, 31-37.
- Fontaine, K. (2005). Management: Arthritis and Health Related Quality of Life. John Hopkins Arthritis Centre.<http://www.hopkins-arthritis.org/patientcorner>.
- Ferrans, C. E., (2007). Differences in what quality-of-life instruments measure NCI monographs, 37, 22-26; doi: 10.1093/jncimonographs/lgm008.
- Ferrans, C.E., Zerwic, J.J., Wilbur, J.E., & Larson, J.L. (2005). Conceptual model of health related quality of life. *Journal of Nursing Scholarship*, 37(4), 336-342. doi: 10.1111/j.1537-5069.2005.00058.
- Fatima, L. (2004) The effect of information booklet provided to caregivers of patients on hemodialysis on knowledge of home care management. *Nursing Journal of India*, Retrieved 14.11.2005.
- Frary, R. (2002). *A Brief Guide to Questionnaire Development*. www.testscoring.vt.edu/fraryquest.html Retrieved on 05.02.2006.
- Farran, C.J., Gilley, D.W., McCann, J.J. & Bienias, J.L. (2007). Efficacy of behavioral intervention for dementia care givers. *Western Journal of Nursing Research*, 29(8), 19442-960. doi:10.1177/0193945907303084. Available from: <http://wjn.sagepub.com>.
- Gordon, H., Guyatt, David H. Feeny, & Donald L. Patrick (1993). Measuring health-related quality of life. *Ann Intern Med*, 118, 622 - 629.
- Gregor, F.M. (2001). Nurses' informal teaching practices: Their nature and impact on the production of patient care. *International Journal of Nursing Studies*, 38, 461-70.
- Grumke, J., & King, K. (1994). Missouri Kidney Program's Patient-Education Program: A 10-Year Review. E-NEPH Archive: *Dialysis & Transplantation*, 23(12), 691-610. Retrieved 05.09.2003.
- Government of India Occupational – educational pattern of employees in India public sector – 1996. Directorate General of Employment & Training. Ministry of Labour, Shram Shakti Bhawan, Rafi Marg, New Delhi 2002.
- Hanly, P. (2007). *Sleep Disorder and End Stage Renal Disease Sleep Medicine Clinics*, 2(1), 59-66.
- Holloway, I., & Wheeler, S., (1996). *Qualitative Research for Nurses*, (1st ed.). Blackwell Science, 39 – 55.

- Himmelfarb, J. (2006). Dialysis at crossroads: Reverse engineering renal replacement therapy, *Cl. J. Am. Soc. Nephrol*, 1,896-902, doi:10.2215/CJN.10911105.
- Heidarnia, A., Dehdan, T., Ghofranipour, F., Kazemnejad, A., & Heidarnia, M. (2009). The effect of health education on health-related quality of life in CABG *Medical Journal of Islamic Republic of Iran* Retrieved: www.cababstractplus.org/abstract.
- Harrison, M B, Lorimer, K., Friedberg, E., Pierscianowski, T. & Brandys. (2005). Leg-ulcer care in the community, before and after implementation of an evidence-based service *CMAJ*. 2005 May 24;172(11):1447-52.
- Hult, H., Lindblad Fridh, M., Lindh Falk, A., & Thörne, K. (2009). Pedagogical processes in healthcare: An exploratory study of pedagogic work with patients and next of kin. *Education for Health*, 22(3), Available from: http://www.educationforhealth.net/
- Harwood, L., Cusolito, H. L., Spittal, J., Wilson, B., & White, S. (2006). Preparing for hemodialysis: patient stressors and responses. *Nephrology Nursing Journal*, 32(3).
- Habibullah, S. (2002). Media health campaign and its impact on knowledge, attitude and practices in the community. *Pak J Med Sci*, 18(3), 232-234.
- Han, S.S., Kim, K.W., Young Na, K., Chae, D. W., Kim, Y.S., Kim, S., & Chin, H.J. (2009). Quality of life and mortality from a nephrologist's view: a prospective observational study. *BMC Nephrology*, 10:39doi:10.1186/1471-2369-10-39 Retrieved from: http://www.biomedcentral.com/1471-2369/10/39
- Higginson, I.J., & Carr, A.J. (2001). Measuring quality of life: using quality of life measures in the clinical setting. *BMJ*, 322: 1927-1300.
- Hays, R.D., & Morales, L.S., (2001). The RAND-36 measure of health-related quality of life. *Ann Med*, 33, 350-357. www.annmed.org Retrieved 19.09.2006.
- Hays, R.D., Kallich, J.D., Mapes, D.L., Coons, S.J., Amin, N., & Carter, W.B. (1995). Kidney disease quality of life short form (KDQOL-SF), Version 1.3: *A Manual for Use and Scoring*. Santa Monica, CA:RAND, P-7994. http://www.rand.org Retrieved 29.12.2006.
- Henrich, W.L., Dialysis as treatment: end stage renal disease 1999 Series editor Schirier R. W. *Atlas of disease of the kidney*, Vol.5, Philadelphia, Blackwell Science.
- Israney, S.M., & Nachane, D.M. (1999). *The Art and Craft of Research Methodologyi In Social Sciences*, (1sted.). Mumbai, Universal Publishing Corporation. 331-362.
- Ira, B., Wilson, & Paul, D. Cleary (1995). Linking clinical variables with health-related quality of life: *A Conceptual Model of Patient Outcomes* *JAMA*, 273, 59 - 65.

- Iseki, K., Kawazoe,N., Osawa, A. & Fukiyama, K. (1993). Survival analysis of dialysis patients in Okinawa, Japan (1971—1990) *Kidney International*; 43, 404-409.
- Jordan, J.E., & Osborne, R.H. (2007). Chronic disease self-management education programs: Challenges ahead. *Medical Journal of Australia*, 18, 84-87.
- Kranse, R.S. (2009). Renal failure chronic and dialysis complications: e Medicine Emergency Medicine, [htt://emedicine.medscape.com](http://emedicine.medscape.com)
- Khanna, U. (2009). The economics of dialysis in India. *Indian Journal of Nephrology*, 19, 1-4.
- Khajehdehi, P., Mojerlar, M., Behzadi, S., & Rais Jalah, E. A. (2001). A randomized double blind placebo controlled trial of supplementing vitamins E C and their combination for treatment of hemodialysis cramps. *Nephrology Dialysis Transplantation*, 16, 1448-1451.
- Kapella, M.C., Lason, J.L., Patel, M.K., Convey, M.K., & Berry, J.K. (2006). Subjective fatigue influencing variables and consequences in chronic obstructive pulmonary disease. *Nursing Research*, 35(1).
- Kallenbach, J.Z., Gutch, C.F., Stoner, M.H. & Corea Anna, L. (2005). *Review of Hemodialysis for Nurses and Dialysis Personnel*, (7th Ed.), St. Louis Elsevier Mosby
- Kalantar-Zadeh, K., Kopple, J.D., Block, G., & Humpreys, M.H., (2001). Association among SF36 quality of life measures and nutrition, hospitalization, and mortality in hemodialysis. *J Am Soc Nephrol* 12, 2797-2806., 2001- p 1-22.
- Kozier B., & Erb G., Berman,A., Snyder, S. (2004). *Fundamentals of Nursing concepts, process and practice*, (7th ed.1st Indian print) Delhi, Saurabh Printers pp 1191-1204
- Kring, D.L. (2008). Using the revised Wilson and Cleary model to explore factors affecting quality of life in persons on hemodialysis. Unpublished Phd Thesis. The University of North Carolina, Greensboro.
- Kidney patient guide. Available form: www.kidneypatientguide.org.uk/SITE/dietfeel.html.
- Kidney School™. A free, online kidney disease learning centre for stages 3, 4 & 5 CKD. Available from: www.kidneyschool.org
- Levey, A.S., Bosh, J.P., Greene, T., & Rogers, N. (1999). Modification of diet in renal disease. *Annals of Internal Medicine* 130 (6). 461 – 470.
- Lukkarman, H., & Hentinen, M. (2006). Treatment of coronary artery disease improve quality of life in the long term. *Nursing Research*.55(1), 26 – 33.

- Lai, C.F., Wang, Y.T., Hung, K.Y., & Peng, Y.S., (2007). Sexual dysfunction in peritoneal dialysis patients. *American Journal of Nephrology*, 27(6), 615 – 621.
- Leggy, V. (2005). Complications of chronic kidney disease. A close look at renal osteodystrophy nutritional disturbances and inflammation. *American Journal of Nursing*, 105(6), 40-50.
- Lee, C.P., Stefanos, A. Z., & Glenn. (2008). M. C. Cost-effectiveness of frequent in-center hemodialysis, *Am Soc Nephrol*. 19(9), 1792–1797. doi: 10.1681/ASN.2008010001.
- Lewis Deborah. (1999). Computer-based approaches to patient education: A review of the literature; JAMIA, 6,272-282 doi:10.1136/jamia.1999.0060272
- Labonte, R. (1994). Health promotion and empowerment. Reflections on professional practice. *Health Education Quarterly*, 21, 253-268.
- Locatelli, F., Fouque, D., Heimburger, O., Drueke, T.B., Cannata-Andia, J.B. Horl, W.H., & Ritz, E. (2002). Nutritional status in dialysis patients: A European consensus. *Nephrology Dialysis Transplantation*, 17,563-572. Retrieved 15-12-2005.
- Lindberg, J., Slowik, M., Figueroa, J.E., Husserl, F., Sadler, R., Savoie, J., Best, J., Szerlip, M., Boudreaus, D., et al (2001). Ochsner Healthy Start Chronic Kidney Disease Clinic: Guide to understanding end stage renal disease. Available from: www.kidney.org
- Meers, C., Hopman, W., Smyu, M.A., Mackenzie, T.A., Mortin, M.A., & Murrary, M. (1995). A Comparison of patient, nurse and physician assessment of health related quality of life in end stage renal disease, *Dialysis and Transplantation*, 24(3), 120.
- Morrin, P.A., (2005). Ethical issues in nephrology. *Journal of Posgrad Med*. 2005 Jul 12, 40:144 – 150.
- Mahajan. B.K &, Gupta, M.C., (1995). *Text Book of Preventive and Social Medicine* (2nded.), 135, New Delhi, Jaypee Brothers Medical Publishers.
- McCarley, P.B. & Salai, P.B., (2005). Cardiovascular diseases in chronic kidney disease recognizing and reducing the risk of common chronic dialysis comorbidity. *American Journal of Nursing*, 105(4).
- Mani, M.K., (2005). Experience with a program for prevention of chronic renal failure in India. *Kidney International*, 67, 975-978; doi:10.1111/j.1523-1755.2005.09419x.
- Mau, I.W., Chiu H.C., Chang P.Y., Hwang S.C., Hwang S.J. (2008). Health related quality of life Taiwanese dialysis patients: Effects of dialysis modality. *The Kahosung Journal of Medical Services*, 24(9), 453-460. Doi:10.1016/s1607-551x(09)7002-6.

- Milazzo, V. (1980). A study of the difference in health knowledge gained through formal and informal teaching. *Heart & Lung*, 9, 1079-1082.
- Mashrabi, O., Rasht, M.D.T., Rasht, M.D.T., Nakhjavani, M.R.J., & Azar, S.A. (2007). Evaluation bone disease in hemodialysis patients. *Research Journal of Medical Sciences* 1 (3), 152-156, 2007.
- Morais, A.A.C., Silva, M.A.T., Faintuch, J., Vidigal, E.J., Costa, R.A., Lyrio, D.C., Trindade, C.R. & Pitanga, K.K. (2005). Correlation of nutritional status and food intake in hemodialysis patients. *Clinics* 60(3) São Paulo. doi: 10.1590/S1807-59322005000300002.
- Mapes, D.L., Lopes, A.A., Satayathum, S., McCullough, K.P., Goodkin, D.A., Locatelli, F., Fukuhara, S., Young, E.W., Kurokawa, K., Saito, A., Bommer, J., Wolfe, R.A., Held, P.J., Port, F.K. (2003). Health-related quality of life as a predictor of mortality and hospitalization: the dialysis outcomes and practice patterns study (DOPPS). *Kidney Int.*, 64(1), 39-49.
- Merkus, M.P., Jager, K.J., Dekker, F.W., de Haan, R.J., Boeschoten, E.W., & Krediet, R.T., (1999). Physical symptoms and quality of life in patients on chronic dialysis: results of The Netherlands cooperative study on adequacy of dialysis (NECOSAD). *Nephrol Dial Transplant*, 14(5), 1163-70.
- Mollaoglu, M. (2004). Depression and health related quality of life in hemodialysis patients. *Dialysis & Transplantation*, 33(9), 544-549.
- Mentari, E.K., DeOreo, P.B., O'Connor, A.S., Love, T.E., Ricanati, E.S., & Sehgal, A.R. (2005). Changes in medicare reimbursement and patient-nephrologist visits, quality of care, and health-related quality of life. *American Journal of Kidney Diseases*, 46(4), 621-627. Retrieved 15.11.07.
- Murphy, S.O., Straub, C.K., & Rosablan, R. (2005). Exercise in the management of fatigue in patients in peritoneal dialysis. *Nephrology Nursing Journal*, 35(5), 468.
- Munro, B.H. (2001). *Statistical methods for health care research*. (4thed.). Philadelphia, Lippincott.
- Munro, B.H., Jacobsen, B.S., Duffy, M.E. (2002). *Introduction to Inferential Statistics and hypothesis testing*. (2nded.), 363 Philadelphia, Lippincott.
- North Carolina Institute of Medicine: (2008). Task force on kidney disease. Addressing chronic kidney disease in North Carolina.
- National Kidney Foundation. (2007). The National Kidney Foundation Kidney Disease Outcomes Quality Initiative. Retrieved on 24.04.07 from <http://www.kidney.org/professionals/KDOQI>.
- Negoeanu, D., Glickman, J.D., & Berns, J.S. (2007). How much is enough? The impact of frequent hemodialysis. *Seminars in dialysis*; 21(2), 192-95.

- Owen, William F., (2003). Jr. Patterns of care for patients with chronic kidney disease in the United States: Dying for improvement. *J Am Soc Nephrol* 14: S76-80.
- Ort, A., Krampitz, S., Pavolich, N. (Eds) (1981). *Research designs*. St. Louis, C.V. Mosby.
- O'Hare, A.M., Choi, A.L., Bertenthal, D., Bachetti, P., Garg, A.V., Kaufman, J.S. et al. (2007). Age affects outcomes in chronic kidney disease. *J Am Soc Nephrol*; 27:2758-2765 doi,10.1681/ASN/2007040422
- Patwardhan, M.B., et al. (2007). Advanced chronic kidney disease practice patterns among nephrologists and non-nephrologists: a database analysis. *Clin J Am Soc Nephrol*, 2(2), 277-83.
- Pezeshki, M.L., & Rostami, Z. (2009). Contributing factors in health-related quality of life. Assessment of ESRD patients: A single center study. *Int J Nephrol Urol*, 1(2),129-136.
- Pagels, A.A., Wang, M., & Wengstrom, Y. (2008). The Impact of a Nurse-Led Clinic on Self-Care Ability, Disease-Specific Knowledge, and Home Dialysis Modality, *Nephrology Nursing Journal*, May-June 2008 Vol. 35, No. 3 May 2008 PP242-243.
- Parker, K.P., Bliwise, D.L., & Rye, D.B. (2000). Hemodialysis disrupts basic sleep regulatory mechanisms: building hypotheses. *Nursing Research*. 49(6), 327-332.
- Pollak, V.F., Lorch, J.A., Shukla, R. & Satwah, S. (2009). The importance of iron in long time survival of hemodialysis patients treated with epoetin-alpha and intravenous iron: Analysis of 9.5 years of prospectively collected data. *B.M.C. Nephrology*.
- Prabahar, M.R., Chandrasekharan, V. & Soundrarajan, P. (2008). Epidemic of chronic kidney disease in India – What can be done? *Saudi J. Kidney transplant*; 19:847-53. Available from <http://www.sjkd.org/text.asp?2008/19/5/847/42478>.
- Phend, C. (2009). Dialysis patients the most pills. *Medpage Today Medicine News: Nephrology, ESRD*. Available from, <http://www.medpagetoday.com/medical-news/VSC>.
- Polit, D.F., & Beck, C.T. (2008). *Nursing Research: Generating and assessing evidence for nursing practice*. (8th ed.). Walter Kluwer, Lippincott.
- Polit, F.B., Hungler, B.P. (1999). *Nursing research, principles and methods*. (5th ed.). Lippincott.
- Promoting excellence: ESRD Table of contents: statistical summary of quality of dialysis patients questionnaires. Promotingexcellence.org

- Quinn, F.M. (1995). *The Principles and practices of nurse education.* (3rd ed.). 113-126, London, Chopman & Hall.
- Quershi, A.R., Alvestrand, A., Danielsson, A., Divino-Filho J.C., Gutierrez, A., Lindholm, B. & Bergstrom, J. (1998). Factors predicting malnutrition in hemodialysis patients; a cross-sectional study. *Kidney International*; 53, 773-82.
- Quality Improvement Collaborative in Asthma Care.* (2005). www.Medscape.com. Retrieved on 12.10.05.
- Questionnaires – the guide to health economics, medical and pharmacy resources. HealthEconomics.Com
- Rosenberg1, M., Kalda1, R., Kasiulevičius, V., & Lember, M. (2007). Management of chronic kidney disease in primary health care: *Position paper of the European Forum of Primary Care.*
- Richard, C.J., (2009). Self-care management in adults undergoing hemodialysis. *Nephrology Nursing Journal.* Available from http://findarticles.com/p/articles/mi_m0ICF/is_4_33/ai_n17214615/
- St Peter, W.L., et al. (2003). Chronic kidney disease: issues and establishing programs and clinics for improved patient outcomes. *Am J Kidney Dis*, 41(5), 903-24.
- Steinberg, E.P. (2003). Improving the quality of care--can we practice what we preach? *N Engl J Med*, 348(26), 2681-3.
- Salisbury, E.M., Game, D.S., Al-Shakarchi, I., Chan, M., Fishman, L., Tookman&, L., Brown, E.A. (2009). Changing practice to improve pain control for renal Patients. pmj.bmjjournals.org Retrieved 23.12.09.
- Shahgholian, N., Ghafourifard, M., Rafieian, M., & Mortazavi, M. (2008). Impact of two types of sodium and ultra filtration profiles on intradialytic hypotension in hemodialysis patients. *Iranian Journal of Nursing and Midwifery Research*, 13(4).
- Song, J.H., Park, G.H., Lee, S.Y., Lee, S.W., Lee, S.W., & Kim, M.J. (2005). Effect of sodium balance and the combination of ultrafiltration profile during sodium profiling hemodialysis on the maintenance of the quality of dialysis and sodium and fluid Balance. *J Am Soc Nephrol* 16, 237-246, doi: 10.1681/ASN.2004070581. Retrieved on 04.01.2006.
- Schloman, B.F., (2004). Health Literacy: A key ingredient for managing personal health. *Online Journal of Issues in Nursing.* <http://nursingworld.org/ojin/infocol/info>. Retrieved on 13.05.2006.
- Simon, S., Stats: So you want to write a questionnaire. www.wlu.ca/lispop/lispop.html. Retrieved on 05.02.2006.

- Smeltzer, S.C. & Bare, B. (20024). *Bruner and Suddarth's text book of medical surgical nursing.* (10th ed.). Philadelphia, Lippincott.
- Sevick, M.A., Trauth, J.M., Ling, B.S., Anderson, R.T., Piatt, G.S., Kibourne, A.M., et al. (2007). Patients with complex chronic diseases: Perspectives on supporting self-management. *J. Gen. Intern M. Suppl* 3,438-444.
- Thomas Hawkens C., & Zazworksy D. (2005). Self management of chronic kidney disease. Patients shoulder the responsibility for day to day management of chronic illness. How can nurses support the autonomy? *American Journal of Nursing*, 105(10), 40-47.
- Tanner, J.L., Craig, C.B., Bartolucci, A.A., Allon, M., Fox, L.M., Geiger, B.F., & Wilson, N.P. (1998). The effect of self monitoring tool on self-efficacy, health beliefs and adherence in patients receiving hemodialysis. *J Ren Nutr.* 8 (4), 203-11.
- Tapiawala S, Vora H, Patel Z, Badve S., & Shah B. (2006). Suggestive global assessment of nutritional status of patients with chronic renal insufficiency and end stage renal disease on dialysis. *JAPI*, 54, 923-26. www.japi.org.
- Tsay S.L., Lee Y.C., & Lee Y.C. (2005). *Issues and innovations in nursing practice.* Effects of an adaptation training program for patient with end stage renal disease: *Journal of Advanced Nursing*. 50(1) 39-46. doi:10.1111/j.1365-2643.2004.03347.
- Tsay S.L., & Hund L. O. (2003). Empowerment of patients with end stage renal stage renal disease – a randomized controlled trial. *International Journal of Nursing Studies*, Vol.1, issue1, January 2004 pp 59-65.
- Tarrass, F., Addou, K., Benjelloun, M., Zamd, M., Medkouri, G., Hachim, K., Benghanem, M.G., & Ramdani, B. (2007). Optimizing hypertension control in hemodialysis patients: a proposed management strategy. *Saudi J Kidney Dis Transpl* [serial online] 18,355-60. Available from: <http://www.sjkdt.org/text.asp?2007/18/3/355/33751>
- Thong1, M.S.Y., Sandra van Dijk, S.V., Noordzij, M., Boeschoten, E.W., Krediet, R.T., Dekker, F.W., & Kaptein, A.A. (2008). for the NECOSAD Symptom clusters in incident dialysis patients: associations with clinical variables and quality of life. *Study Group Nephrol Dial Transplant* (2008) 1–6. doi: 10.1093/ndt/gfn449.
- Turner, S., Wellard, S., & Bethune, E. (1999). Registered nurses' perception of teaching. Constraints to the teaching moment. *International Journal of Nursing Practice*, 5,14-20.
- Thomas, C.L. (1988). Edited by Taber's cyclopedic *medical dictionary*, New Delhi, Jaypee brothers. The 5-Item Version of the International Index of Erectile Dysfunction (IIEF-5) Institute for Algorithmic Medicine.

The 5-item version of the international index of erectile dysfunction (IIEF-5) – Institute of Algorithmic Medicine. Retrieved on 11.10.2006.

Unruh, M.L., Busy bee, D.J., Dew, M.A., Evan, I., Wu, W.A., Fink, N.E. et.al. (2006). Sleep quality and its correlated in the first year of dialysis. *Clin. J. Am. Soc. Nephrol.*, 1:802-810 doi: 10.2215/C. J. N./.00710-206.

Unruh, M., Yan, G., Radeva, M., Hays, R.D., Benz, R., Athienites, N.V., Kusek, J., Levey, A.S., & Meyer, K.B., and the HEMO Study Group (2003) - Bias in assessment of health-related quality of life in a hemodialysis population: A comparison of self-administered and interviewer-administered surveys in the hemo study *J Am Soc Nephrol* 14, 2132-2141, *American Society of Nephrology*.

Unruh, M., Miskulin, D., Yan, G., Hays, R.D., Benz, R., John W. Kusek., & J.W. And Meyer, K.B., The Hemo Study Group. (2004). Racial differences in health-related quality of life among hemodialysis patients. *Kidney International* 65, 1482–1491; doi:10.1111/j.1523-1755.2004.00529.x

Vijaykumar, K., & Varghese, R.T. (2009). Quality of life among diabetic subjects: Indian perspectives. Hand book of disease burdens and quality of life measures. 2071-2093: Springer New York. doi:10.1007/978-0-387-78665-0_121. Retrieved 30, 09. 2009.

Wingard. R.L, Chan, K.E., Lazarus, J. M., & Hakim, R. M., (2009). The "Right" of Passage: Surviving the first year of dialysis *Clin. J. Am. Soc. Nephrol.*, 4(Supplement_1): S114 - S120.

Wilson, I.B., & Cleary, P.D. (1995). Linking clinical variables with health-related quality of life: a conceptual model of patient outcomes. *J Am Med Assoc.* 1995; 273:59–65.

Weisbord, S. D., Fried, L.F., Mor, M.K., Rensuk, A.L., & Unruh, M.L., (2007). Renal provider recognition of symptoms in patients on maintenance hemodialysis. *Clinical Journal American Society of Nephrology* 2(5), 960-967.

www.hemodialysis.org. (2009). Dialysis and your (health related) quality of Life. *Medical Education Institute*.

Wells, C. (2003). Optimizing nutrition in patients with chronic kidney disease *Nephrology Nursing Journal*. 30(6), 637-648.

Wojciechowski & Cichowski, K. (2007). A case review; designing new patient education system. *The Internet Journal of Advanced Nursing Practice*. 8(2), ISSN 1523-6064.

Walsh M., Culleton B., Tonelli M., & Manns B. (2005). A systematic review of the effect of nocturnal hemodialysis on blood pressure, left ventricular hypertrophy, anemia mineral metabolism and health related quality of life. *Kidney International* 67, 1500.1508. doi:10:1111/j.1523-1755:2005.00228.

- Wingard, R.L., Pupim, L.B., Krishnan, M., Shintani, A., Ikizler, T.A., & Hakim, R.M. (2007). Early intervention improves mortality and hospitalization rates in incident hemodialysis patients: Right start program *Clin J Am Soc Nephrol.* 2(6), 1170-5.
- Wikstrom, Bjom, (2007). Itchy skin - a clinical problem for haemodialysis patients. *Nephrol Dial Transplant*, 22 [Suppl 5]: v3-v7.
- Weisbord, S.D., Fried, L.F., Arnold, R.M., Fine, M.J., Levenson, D.J., Peterson, & R.A., Switzer, G.E. (2005). Prevalence, severity, and importance of physical and emotional symptoms in chronic hemodialysis patients. *J. Am. Soc. Nephrol.* 16(8), 2487-94. Epub 2005 Jun 23. PMID: 15975996
- Weisbord, S.D., Fried, L.F., Arnold, R.M., Rotondi, A.J., Fine, M.J., Levenson, D.J. et al. (2004). Development of a symptom assessment instrument for chronic hemodialysis patients: The dialysis symptom index. *J. pain symptom manage*; 27(3): 226 to 240. doi:10.1016/j.painsymman.2003.07.004
- White, A.C., Pilkey, R.M., Lam, M., & Holland, D.C., (2002). Pre-dialysis clinic attendance improves quality of life among hemodialysis patients. *BMC Nephrol.* 3,3, Published online. Retrieved 30.09.05.
- Wu, A.W., Fink, N.E., Cagney, K.A., Bass, E.B., Rubin, H.R., Meyer, K.B., Sadler, J.H., & Powe, N.R. (2001). Developing a Health-related quality of life Measure for End-Stage Renal Disease: The CHOICE Health experience Questionnaire. *American Journal of Kidney Diseases*, Vol.37, No.1 (January), 2001: pp 11-21.
- Ware, J.E., SF36 Health Survey Update. <http://www.sf-36.com/> and <http://www.qualitymetric.com/>
- Weigel, M., Meston, C., & Rosen, R., (2006). The female sexual function index (FSFI): cross-validation and development of clinical cutoff scores. *J Sex Marital Ther*, 31(1),1-20. PubMed. Retrieved 15.12.2006.
- World kidney day. (2008). <http://www.worldkidneyday.org>
- W.H.O. Quality of life-BREF (WHOQOL-BREF) World Health Organization 2005. Retrieved 14.07.2005.
- Wilson, I. B., & Cleary, P. D. (1995). Linking clinical variables with health-related quality of life. *Journal of the American Medical Association*, 273, 59-65.
- Yang, S.C., Kuo, P.W., Wnag, J.D., Lin, M.I. & Su, S. (2005). Quality of life and its determinants of hemodialysis patients in Taiwan measured with WHOQOL-BREF (TW). *American Journal of Kidney diseases*, 46(4), 635-641.